

## Cyrex Array 10 Sample Report

TEST		RESULT		
Array 10 - Food Immune Reactivity Screen **	IN RANGE (Normal)	EQUIVOCAL*	OUT OF RANGE	REFERENCE (ELISA Index)
DAIRY and EGGS, Modified				
Egg White, cooked	0.75			0.1-1.6
Egg Yolk, cooked	0.85			0.1-1.7
Goat's Milk	0.70			0.1-1.9
Soft Cheese + Hard Cheese	0.54			0.1-1.7
Yogurt	0.74			0.1-2.0
GRAINS, Raw and Modified				
Rice, white + brown, cooked	0.48			0.1-1.3
Rice Cake	0.42			0.2-1.8
Rice Protein	0.81			0.2-1.7
Rice Endochitinase	1.08			0.2-1.7
Wild Rice, cooked	0.88			0.1-1.3
Wheat + Alpha-Gliadins	0.73			0.2-1.9
BEANS and LEGUMES, Modified				
Black Bean, cooked	0.26			0.3-2.1
Bean Agglutinins	0.90			0.3-1.9
Dark Chocolate + Cocoa	0.40			0.2-1.2
Fava Bean, cooked		1.20		0.3-1.5
Garbanzo Bean, cooked	0.58			0.2-1.8
Kidney Bean, cooked	0.39			0.3-1.5
Lentil, cooked	1.27			0.3-2.0
Lentil Lectin	1.26			0.2-1.9
Lima Bean, cooked	0.82			0.1-1.8
Pinto Bean, cooked	0.36			0.4-2.4
Soybean Agglutinin	0.64			0.1-1.7
Soybean Oleosin + Aquaporin	0.91			0.2-1.8
Soy Sauce, gluten-free	0.71			0.2-1.9

<sup>\*\*</sup> For details on the method of cooking, please see specification sheets. All analytes are tested for IgG and IgA combined.

Mark G. Kartub, M.D., Medical Director

<sup>\*</sup> Reference ranges are calculated based on the mean ±2 standard deviations (SD). Results > 1 SD, and <2 SDs above the mean are considered to be equivocal. An equivocal result represents the range between negative and suspicious low positive results. Results >2 SDs are considered out of range, and positive.



Tofu	1.04		0.2-1.4
NUTS and SEEDS, Raw and Modified			
Almond	0.78		0.2-1.8
Almond, roasted	0.63		0.2-2.0
Brazil Nut, raw + roasted	0.63		0.1-1.8
Cashew	0.79		0.2-1.5
Cashew, roasted	1.11		0.2-2.3
Cashew Vicilin	0.65		0.3-1.7
Chia Seed	0.46		0.2-1.7
Flax Seed	0.87		0.1-1.3
Hazelnut, raw + roasted	0.94		0.1-1.7
Macadamia Nut, raw + roasted	1.27		0.3-2.3
Mustard Seed		1.46	0.4-1.5
Pecan, raw + roasted	0.91		0.3-1.5
Peanut, roasted	0.40		0.2-1.4
Peanut Butter	0.70		0.2-1.9
Peanut Agglutinin	1.29		0.3-1.9
Peanut Oleosin	1.13		0.3-1.8
Pistachio, raw + roasted	0.75		0.4-2.0
Pumpkin Seeds, roasted	0.78		0.2-1.6
Sesame Albumin	0.66		0.2-1.3
Sesame Oleosin	0.61		0.2-1.6
Sunflower Seeds, roasted	0.53		0.2-1.5
Walnut	1.55		0.3-2.0
VEGETABLES, Raw and Modified			
Artichoke, cooked	0.88		0.1-2.7
Asparagus	0.82		0.3-2.1
Asparagus, cooked	0.61		0.1-2.2
Beet, cooked	0.46		0.1-1.5
Bell Pepper		1.55	0.1-1.8
Broccoli	0.89		0.1-1.5

<sup>\*\*</sup> For details on the method of cooking, please see specification sheets. All analytes are tested for IgG and IgA combined.

Mark G. Kartub, M.D., Medical Director

<sup>\*</sup> Reference ranges are calculated based on the mean ±2 standard deviations (SD). Results > 1 SD, and <2 SDs above the mean are considered to be equivocal. An equivocal result represents the range between negative and suspicious low positive results. Results >2 SDs are considered out of range, and positive.



Broccoli, cooked	0.58		0.1-2.0
Brussels Sprouts, cooked	0.54		0.1-3.0
Cabbage, red + green	0.92		0.1-2.5
Cabbage, red + green, cooked	0.62		0.1-2.5
Canola Oleosin	0.87		0.1-1.9
Carrot	0.59		0.1-2.7
Carrot, cooked	0.49		0.1-2.2
Cauliflower, cooked	0.44		0.1-2.2
Celery	0.76		0.1-2.3
Chili Pepper	1.21		0.1-1.9
Corn + Aquaporin, cooked	0.55		0.1-1.8
Popped Corn	0.71		0.1-1.9
Corn Oleosin	0.19		0.1-1.4
Cucumber, pickled	0.51		0.1-2.6
Eggplant, cooked	1.07		0.1-2.1
Garlic	0.44		0.1-2.2
Garlic, cooked	0.53		0.1-1.9
Green Bean, cooked	0.45		0.1-1.5
Lettuce	0.51		0.1-1.5
Mushroom, raw + cooked	0.75		0.1-1.6
Okra, cooked	0.63		0.1-1.5
Olive, green + black, pickled	0.88		0.1-1.7
Onion + Scallion	0.67		0.1-1.7
Onion + Scallion, cooked	0.24		0.1-1.5
Pea, cooked	0.71		0.1-1.5
Pea Protein		3.12	0.1-2.3
Pea Lectin		1.74	0.1-1.7
Potato, white, cooked (baked)	0.34		0.1-1.8
Potato, white, cooked (fried)	0.86		0.1-1.6
Pumpkin + Squash, cooked		1.01	0.1-1.3
Radish		1.64	0.1-1.7

<sup>\*\*</sup> For details on the method of cooking, please see specification sheets. All analytes are tested for IgG and IgA combined.

Mark G. Kartub, M.D., Medical Director

<sup>\*</sup> Reference ranges are calculated based on the mean ±2 standard deviations (SD). Results > 1 SD, and <2 SDs above the mean are considered to be equivocal. An equivocal result represents the range between negative and suspicious low positive results. Results >2 SDs are considered out of range, and positive.



Safflower + Sunflower Oleosin	0.55		0.1-1.5
Seaweed		0.96	0.1-1.2
Spinach + Aquaporin	0.74		0.1-1.5
Tomato + Aquaporin	1.00		0.2-2.2
Tomato Paste	0.59		0.2-2.1
Yam + Sweet Potato, cooked	0.73		0.3-1.9
Zucchini, cooked	0.53		0.3-1.9
FRUIT, Raw and Modified			
Apple	0.88		0.2-1.5
Apple Cider	0.80		0.3-1.3
Apricot	1.58		0.2-2.8
Avocado	1.30		0.6-2.5
Banana	0.17		0.1-2.3
Banana, cooked	0.26		0.2-2.8
Latex Hevein	0.33		0.3-2.0
Blueberry	0.79		0.1-1.6
Cantaloupe + Honeydew Melon	0.50		0.1-1.2
Cherry		1.36	0.2-1.4
Coconut, meat + water	0.92		0.2-2.0
Cranberry	0.84		0.3-2.4
Date	0.66		0.2-1.4
Fig	0.85		0.2-2.2
Grape, red + green	0.60		0.2-1.0
Red Wine	1.21		0.1-2.3
White Wine		1.89	0.1-2.6
Grapefruit	0.80		0.2-1.9
Kiwi	0.56		0.2-1.7
Lemon + Lime	0.43		0.2-1.3
Mango	0.48		0.2-1.5
Orange	1.14		0.2-1.7
Orange Juice	0.65		0.2-1.8

<sup>\*\*</sup> For details on the method of cooking, please see specification sheets. All analytes are tested for IgG and IgA combined.

Mark G. Kartub, M.D., Medical Director

<sup>\*</sup> Reference ranges are calculated based on the mean ±2 standard deviations (SD). Results > 1 SD, and <2 SDs above the mean are considered to be equivocal. An equivocal result represents the range between negative and suspicious low positive results. Results >2 SDs are considered out of range, and positive.



Papaya	1.11		0.2-1.7
Peach + Nectarine		2.30	0.2-2.0
Pear	0.63		0.2-2.6
Pineapple	0.35		0.1-1.9
Pineapple Bromelain	0.84		0.2-2.6
Plum	0.84		0.3-2.2
Pomegranate	0.97		0.4-2.2
Strawberry	0.84		0.3-2.3
Watermelon	0.55		0.2-1.8
FISH and SEAFOOD, Raw and Modified			
Cod, cooked	1.25		0.2-1.8
Halibut, cooked	1.04		0.1-1.6
Mackerel, cooked	0.94		0.2-2.0
Red Snapper, cooked	0.98		0.1-1.5
Salmon	0.74		0.2-2.3
Salmon, cooked	0.38		0.2-2.4
Sardine + Anchovy, cooked	0.67		0.3-2.9
Sea Bass, cooked	0.53		0.2-2.8
Tilapia, cooked	0.79		0.1-1.8
Trout, cooked	0.81		0.1-2.4
Tuna	0.50		0.1-2.7
Tuna, cooked	0.39		0.1-1.3
Whitefish, cooked	0.51		0.1-1.4
Crab + Lobster, cooked	0.82		0.2-2.1
Imitation Crab, cooked	0.65		0.1-1.7
Clam, cooked	0.78		0.1-1.9
Oyster, cooked	0.84		0.1-1.6
Scallops, cooked	0.81		0.1-2.0
Squid (Calamari), cooked	0.85		0.1-2.0
Shrimp, cooked	0.48		0.1-2.1
Shrimp Tropomyosin	0.91		0.1-1.6

<sup>\*\*</sup> For details on the method of cooking, please see specification sheets. All analytes are tested for IgG and IgA combined.

Mark G. Kartub, M.D., Medical Director

<sup>\*</sup> Reference ranges are calculated based on the mean ±2 standard deviations (SD). Results > 1 SD, and <2 SDs above the mean are considered to be equivocal. An equivocal result represents the range between negative and suspicious low positive results. Results >2 SDs are considered out of range, and positive.



Parvalbumin	0.95			0.1-1.7
MEAT, Modified				
Beef, cooked medium	0.79			0.3-1.9
Chicken, cooked	1.08			0.2-1.5
Lamb, cooked	0.57			0.1-1.3
Pork, cooked	0.28			0.1-2.2
Turkey, cooked	0.19			0.1-1.3
Gelatin	0.48			0.1-1.3
Meat Glue	0.40			0.1-1.3
HERBS, Raw				
Basil			1.87	0.2-1.8
Cilantro	0.83			0.1-1.5
Cumin			>3.00	0.2-2.3
Dill	0.88			0.3-1.7
Mint			2.36	0.3-2.1
Oregano	0.60			0.4-2.6
Parsley	0.51			0.1-1.3
Rosemary	1.02			0.3-2.2
Thyme			2.21	0.4-1.8
SPICES, Raw				
Cinnamon	0.79			0.3-1.7
Clove	0.86			0.4-1.8
Ginger	0.86			0.1-2.5
Nutmeg	0.87			0.2-1.9
Paprika		2.08		0.2-2.1
Turmeric (Curcumin)	0.40			0.1-1.7
Vanilla	0.94			0.1-2.4
GUMS				
Beta-Glucan	0.37			0.1-1.3
Carrageenan	0.58			0.2-2.0
Gum Guar	0.56			0.2-2.4

<sup>\*\*</sup> For details on the method of cooking, please see specification sheets. All analytes are tested for IgG and IgA combined.

Mark G. Kartub, M.D., Medical Director

<sup>\*</sup> Reference ranges are calculated based on the mean ±2 standard deviations (SD). Results > 1 SD, and <2 SDs above the mean are considered to be equivocal. An equivocal result represents the range between negative and suspicious low positive results. Results >2 SDs are considered out of range, and positive.



Gum Tragacanth	0.20		0.1-1.4
Locust Bean Gum	0.44		0.2-1.4
Mastic Gum + Gum Arabic	0.32		0.1-1.1
Xanthan Gum	0.58		0.1-1.7
BREWED BEVERAGES and ADDITIVES			
Coffee Bean Protein, brewed	0.41		0.2-1.8
Black Tea, brewed	0.96		0.3-1.6
Green Tea, brewed	0.83		0.3-1.8
Honey, raw +processed		0.99	0.1-1.3
Food Coloring	0.67		0.2-1.8

Mark G. Kartub, M.D., Medical Director

<sup>\*\*</sup> For details on the method of cooking, please see specification sheets. All analytes are tested for IgG and IgA combined.

<sup>\*</sup> Reference ranges are calculated based on the mean ±2 standard deviations (SD). Results > 1 SD, and <2 SDs above the mean are considered to be equivocal. An equivocal result represents the range between negative and suspicious low positive results. Results >2 SDs are considered out of range, and positive.